

NEUPRO® (rotigotine transdermal system) is a prescription medicine used to treat moderate-to-severe primary Restless Legs Syndrome.

These are legs.
They help me be active.
I just wish they would
let me rest.

This is NEUPRO for RLS.

Eligible commercial patients may pay as little
as \$10 per 30-day supply of NEUPRO*

*Eligibility terms and conditions apply.

When Restless Legs Syndrome (RLS)
interferes with your life, it's
time to talk about NEUPRO

 **Neupro®**
(rotigotine transdermal system)
www.NEUPRO.com

It's Time to Talk About NEUPRO.

The
24HR
PATCH

Please see Important Safety Information within and refer to Patient Information for NEUPRO at the end of this brochure or at www.NEUPRO.com.


Congratulations, you've taken a step in the right direction. Talking to your doctor about your Restless Legs Syndrome (RLS), also known as Willis-Ekbom disease, is essential for managing your symptoms. There's a reason your doctor chose to prescribe NEUPRO® (rotigotine transdermal system) for you. NEUPRO is a medication with the proven potential for RLS symptom relief. Patients with moderate-to-severe primary RLS showed improvement in RLS symptoms at the end of a 6-month clinical study. In addition, nearly 1 out of 4 patients (23%) taking NEUPRO were completely symptom-free at the end of the study, compared to nearly 1 out of 10 patients (9%) who used a placebo patch, without any medicine. The most common side effects in people taking NEUPRO for RLS are application site reactions, nausea, difficulty falling asleep and staying asleep, sleepiness, and headache.




Actual patch.
Not shown actual size.

Important Safety Information

NEUPRO contains a sulfite called sodium metabisulfite. Sulfites can cause severe allergic reactions that are life threatening to some people who are sensitive to sulfites. People with asthma are more sensitive to sulfites. Remove the patch right away and call your doctor if you have swelling of the lips or tongue, chest pain, trouble breathing or swallowing.



These are legs.
They can take me on a bike ride.
I wish they would let me take a long car ride.



What Is RLS?

Restless Legs Syndrome (RLS)* is a neurological disorder that results in a powerful urge to move your legs (and, sometimes, other parts of your body).

RLS affects up to 23 million American adults; it is estimated that a third have moderate-to-severe RLS. Symptoms tend to get worse in the evening and at night, and keep people from resting and relaxing, and may even affect sleep.

*Restless Legs Syndrome is also known as Willis-Ekbom disease.

Important Safety Information

NEUPRO may make you fall asleep suddenly or without warning while doing normal activities, such as driving, which may result in accidents.

Tell your doctor right away if this happens. Drinking alcohol or taking other medicines that cause drowsiness may increase your chances of becoming sleepy while using NEUPRO. Do not drive, use hazardous machinery, or do other dangerous activities until you know how NEUPRO affects you.

Doctors use four key points—sometimes called the “**URGE**” criteria—to diagnose RLS:


Urge to move the legs—sometimes accompanied by uncomfortable sensations deep in the legs that may be described as tingling, creeping, crawling, itching, or burning.

Rest-induced — the urge to move gets worse during inactivity — resting, sitting, or lying down.

Gets better with activity — movement such as walking or stretching brings relief, but unpleasant sensations reappear when you stop.

Evening and night—the urge to move increases in the evening or at night or occurs only in the evening or at night.

Please see Important Safety Information within and refer to Patient Information for NEUPRO at the end of this brochure or at www.NEUPRO.com.



These are legs.
They keep me moving.
Even when I am trying to sleep.



Why NEUPRO May Be Right for You

NEUPRO® (rotigotine transdermal system) is a prescription medicine used to treat moderate-to-severe primary Restless Legs Syndrome. It's a patch that delivers the dopamine agonist rotigotine through the skin. With NEUPRO, your RLS symptoms are treated with one patch, once a day.

In a 6-month clinical trial, patients with moderate-to-severe primary RLS taking NEUPRO saw improvement in RLS symptoms, as measured by a combined score based on the following 10 items:

1. RLS discomfort in arms and legs
2. Need to move around because of RLS symptoms
3. Relief of RLS discomfort from moving around
4. Sleep disturbance due to RLS symptoms
5. Tiredness or sleepiness during the day due to RLS symptoms
6. Severity of RLS as a whole
7. Frequency of RLS symptoms
8. Severity of RLS symptoms
9. Impact of RLS symptoms on daily activities
10. Severity of mood disturbance due to RLS symptoms

In fact, nearly 1 out of 4 patients (23%) taking NEUPRO were completely symptom free at the end of the study (compared to nearly 1 out of 10 patients (9%) who used a placebo patch, without any medicine). In other words, these patients had a score of zero on all 10 items, which means no symptoms.

Do not use NEUPRO if you are allergic to rotigotine or any of the other ingredients in the patch.

Important Safety Information

NEUPRO can cause decreases in blood pressure, especially when you start or increase your dose. Increases in blood pressure and heart rate, and fainting, also can occur. If you faint or feel dizzy, nauseated, or sweaty when you stand up from sitting or lying down, tell your doctor.

Skin reactions may occur at the site where you apply NEUPRO. Tell your doctor if you get a rash, redness, swelling, or itching that will not go away.

Please see Important Safety Information within and refer to Patient Information for NEUPRO at the end of this brochure or at www.NEUPRO.com.

NEUPRO is
a once a day
patch that
provides
medicine
for 24 hours.



These are legs.

They let me walk to the mailbox.
I wish they would let me fly to visit family.



Why a Patch May Be Right for You

- NEUPRO® (rotigotine transdermal system) is a patch that offers once-daily dosing. Simply choose the time of day that works best for you and apply the patch at the same time each day. There's no need to schedule NEUPRO around bedtime.
- NEUPRO provides stable, continuous delivery 24 hours a day, so you get a consistent level of medication throughout the day.
- The medication in the NEUPRO Patch does not go through your digestive system
 - Food won't interfere with how it works
 - No pills to swallow
 - No need to schedule NEUPRO around meals
 - The medication in NEUPRO will not be affected by the digestive tract

NEUPRO can cause nausea, vomiting, indigestion, or upset stomach, which may occur more frequently when you first start using the patch.

Skin reactions may occur at the site where you apply NEUPRO. Tell your doctor if you get a rash, redness, swelling, or itching that will not go away.


Important Safety Information

NEUPRO can cause psychotic symptoms including hallucinations (seeing or hearing things that are not real), confusion, excessive suspicion, aggressive behavior, agitation, delusional beliefs (believing things that are not real), and disorganized thinking. If you have hallucinations or any of these other psychotic-like changes, talk with your doctor right away.

Some patients using NEUPRO get urges to behave in a way that is unusual for them, such as unusual urges to gamble, strong urges to spend money, binge eating, or increased sexual urges and behaviors. Some patients may want to use more NEUPRO than prescribed for their symptoms (dopamine dysregulation syndrome). If you or your family notices you are developing any unusual behaviors, talk to your doctor.

NEUPRO may cause Restless Legs Syndrome symptoms to come back (rebound), become worse, or start earlier in the day.

Please see Important Safety Information within and refer to Patient Information for NEUPRO at the end of this brochure or at www.NEUPRO.com.



These are legs.
They keep me moving.
Even when I don't want them to.

Applying the Patch

It is important to apply the patch correctly.



Visit www.NEUPRO.com to watch the "How to Apply the Patch" video for complete step-by-step instructions. Be sure to ask your doctor or pharmacist about anything you do not understand.

Frequently Asked Questions About NEUPRO

1. Can I get my NEUPRO Patch wet?

Yes. You can bathe, shower, or swim while wearing the NEUPRO Patch. However, water may loosen your patch. Avoid hot baths, as the heat may cause too much medicine to pass through the skin.

2. What if my NEUPRO Patch falls off?

If your patch falls off, put on a fresh patch until the time you usually change your patch. Then put a new patch on a different part of your skin at your regular scheduled time. If the edges of the patch lift, you may tape them down with bandaging tape.

3. What if I forget to put on my NEUPRO Patch?

If you miss a dose or forget to change your NEUPRO patch, apply a new NEUPRO patch as soon as you remember. Replace the NEUPRO patch at your normal time the next day.


4. How long does it take for the NEUPRO Patch to start working?

Your doctor should start you on a low dose of NEUPRO. Your doctor may change the dose weekly until you are taking the right amount of medicine to control your symptoms. It may take several weeks before you reach the dose that controls your symptoms best.

Important Safety Information


Avoid exposing the NEUPRO Patch you are wearing to heating pads, electric blankets, heat lamps, saunas, hot tubs, heated water beds, and direct sunlight. Too much medicine could be absorbed into your body. Also, do not wear NEUPRO during medical procedures called magnetic resonance imaging (MRI) or cardioversion because this could cause skin burns.

Please see Important Safety Information within and refer to Patient Information for NEUPRO at the end of this brochure or at www.NEUPRO.com.



These are legs.

They aren't just any legs. They're my legs. I rely on them...and I need them to let me rest.



Frequently Asked Questions, continued

5. What should I avoid while taking NEUPRO?

Do not drive, operate machinery, or do other dangerous activities until you know how NEUPRO affects you.

Avoid exposing the NEUPRO Patch you are wearing to heating pads, electric blankets, heat lamps, saunas, hot tubs, heated waterbeds, and direct sunlight. Too much medicine could be absorbed into your body.

Do not use NEUPRO during certain medical procedures called magnetic resonance imaging (MRI) or cardioversion. Using NEUPRO during these procedures could cause a burn to the site where you applied your NEUPRO Patch.

6. Do I have to apply my NEUPRO Patch around mealtime?

No. Apply your patch when it's convenient for you, and apply it at the same time each day. Because NEUPRO is delivered through the skin, food should not affect the medication.

7. Why do I need to hold the NEUPRO Patch on for at least 30 seconds?

Press your NEUPRO Patch firmly with the palm of your hand for 30 seconds to make sure there is good contact with your skin, especially around the edges. The warmth of your hand helps the adhesive on the patch stick to your skin. Make sure that your NEUPRO Patch is flat against your skin. There should be no bumps or folds in your NEUPRO Patch.

8. How can I prevent skin irritation or other skin-related side effects?

You need to change the location of the NEUPRO Patch on a daily basis and should not use the same location for 14 days. Changing the location will reduce the chance of skin irritation. You may see mild redness, like when you remove an adhesive bandage, which is normal. Of course, always check with your doctor if you have any concerns.

Patients should report application site reactions that do not go away after a few days, that get worse, or that spread outside the patch site. If there is a skin rash or irritation from the patch, direct sunlight on the area should be avoided until the skin heals. Exposure could lead to changes in skin color.

The NEUPRO Patch Placement Tracker can help you rotate your application site every day. Visit www.NEUPRO.com to download the Tracker now.

9. How do I safely stop using the NEUPRO Patch?

Do not stop using NEUPRO without talking to your doctor first. If your doctor tells you to stop using NEUPRO, you should ask your doctor for specific instructions on how to slowly and safely discontinue using NEUPRO. If you stop using NEUPRO, you may have withdrawal symptoms.

10. What does the NEUPRO Patch look like?

NEUPRO® (rotigotine transdermal system) is a small, thin patch that comes in 3 sizes based on dosage, 1 mg, 2 mg, and 3 mg.


11. How can I save on NEUPRO?

The NEUPRO Patient Savings Program™ is here to help.

Visit www.NEUPRO.com to see if you are eligible to save today.*

*Eligibility terms and conditions apply.

Call [ucbCARES® at 1-844-599-2273](tel:1-844-599-2273) if you have questions about how to apply the NEUPRO Patch.



Actual patches. Not shown actual size.

Important Safety Information

Do not stop using NEUPRO without talking to your doctor first. If your doctor tells you to stop using NEUPRO, you should ask your doctor for specific instructions on how to slowly and safely discontinue using NEUPRO. If you stop using NEUPRO, you may have withdrawal symptoms such as fever, confusion, severe muscle stiffness, feeling like you do not care about the things you usually care about (apathy), anxiety, depression, fatigue, insomnia, sweating and pain.

Tell your doctor if you have breathing problems, a sleep disorder, mental problems, high or low blood pressure, heart problems; are pregnant or plan to become pregnant; or are breastfeeding or plan to breastfeed. NEUPRO may not be right for you.

The most common side effects in people taking NEUPRO for RLS are application site reactions, nausea, difficulty falling asleep and staying asleep, sleepiness, and headache.

These are not all the possible side effects of NEUPRO. For more information, ask your doctor or pharmacist.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to UCB, Inc. at [ucbCARES® \(1-844-599-2273\)](tel:1-844-599-2273).

Please see accompanying Patient Information about the NEUPRO Patch. This information does not take the place of talking with your healthcare provider about your condition or treatment.

Please see Important Safety Information within and refer to Patient Information for NEUPRO at the end of this brochure or at www.NEUPRO.com.

Important Safety Information

NEUPRO contains a sulfite called sodium metabisulfite. Sulfites can cause severe allergic reactions that are life threatening to some people who are sensitive to sulfites. People with asthma are more sensitive to sulfites. Remove the patch right away and call your doctor if you have swelling of the lips or tongue, chest pain, or trouble breathing or swallowing.

NEUPRO may make you fall asleep suddenly or without warning while doing normal activities, such as driving, which may result in accidents. Tell your doctor right away if this happens. Drinking alcohol or taking other medicines that cause drowsiness may increase your chances of becoming sleepy while using NEUPRO. Do not drive, use hazardous machinery, or do other dangerous activities until you know how NEUPRO affects you.

NEUPRO can cause psychotic symptoms including hallucinations (seeing or hearing things that are not real), confusion, excessive suspicion, aggressive behavior, agitation, delusional beliefs (believing things that are not real), and disorganized thinking. If you have hallucinations or any of these other psychotic-like changes, talk with your doctor right away.

NEUPRO can cause decreases in blood pressure, especially when you start or increase your dose. Increases in blood pressure and heart rate, and fainting, also can occur. If you faint or feel dizzy, nauseated, or sweaty when you stand up from sitting or lying down, tell your doctor.

Some patients using NEUPRO get urges to behave in a way that is unusual for them, such as unusual urges to gamble, strong urges to spend money, binge eating, or increased sexual urges and behaviors. Some patients may want to use more NEUPRO than prescribed for their symptoms (dopamine dysregulation syndrome). If you or your family notices you are developing any unusual behaviors, talk to your doctor.

NEUPRO may cause Restless Legs Syndrome symptoms to come back (rebound), become worse, or start earlier in the day.

Skin reactions may occur at the site where you apply NEUPRO. Tell your doctor if you get a rash, redness, swelling, or itching that will not go away.

Avoid exposing the NEUPRO patch you are wearing to heating pads, electric blankets, heat lamps, saunas, hot tubs, heated water beds, and direct sunlight. Too much medicine could be absorbed into your body. Also, do not wear NEUPRO during medical procedures called magnetic resonance imaging (MRI) or cardioversion because this could cause skin burns.

Do not stop using NEUPRO without talking to your doctor first. If your doctor tells you to stop using NEUPRO, you should ask your doctor for specific instructions on how to slowly and safely discontinue using NEUPRO. If you stop using NEUPRO, you may have withdrawal symptoms such as fever, confusion, severe muscle stiffness, feeling like you do not care about the things you usually care about (apathy), anxiety, depression, fatigue, insomnia, sweating and pain.

Tell your doctor if you have breathing problems, a sleep disorder, mental problems, high or low blood pressure, or heart problems; are pregnant or plan to become pregnant; or are breastfeeding or plan to breastfeed. NEUPRO may not be right for you.

The most common side effects in patients taking NEUPRO for Restless Legs Syndrome are application site reactions, nausea, difficulty falling asleep and staying asleep, sleepiness, and headache.

These are not all the possible side effects of NEUPRO. For more information, ask your doctor or pharmacist.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report side effects to UCB, Inc. at ucbCARES.com (1-844-599-2273).

Please see accompanying Patient Information about the NEUPRO Patch. This information does not take the place of talking with your healthcare provider about your condition or treatment.


Please see Important Safety Information throughout and refer to the Patient Information at the end of this brochure or at www.NEUPRO.com.

 **Neupro**[®]
(rotigotine transdermal system)

These are legs.
This is NEUPRO for RLS.

The
24HR
PATCH

- 24 hours of continuous medication
- Once-daily dosing
- Proven potential for complete symptom relief in moderate-to-severe primary RLS



Savings Offer

Eligible commercial patients may pay as little as \$10 per 30-day supply of NEUPRO.*

*Eligibility terms and conditions apply.

Learn more and sign up for the NEUPRO Patient Savings Program™ at www.NEUPRO.com

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Important Safety Information

NEUPRO can cause serious side effects including severe allergic reactions to the ingredient sodium metabisulfite, falling asleep during normal activities, hallucinations and other psychosis, increases or decreases in blood pressure and heart rate, fainting, unusual urges, skin site reactions, RLS symptoms that return, worsen or start earlier in the day, and withdrawal symptoms as the dose is slowly lowered or stopped.

 **Neupro**[®]
(rotigotine transdermal system)



NEUPRO[®] and ucBCARES[®] are registered trademarks, and the NEUPRO Patient Savings Program™ is a trademark, of the UCB Group of Companies.
©2021 UCB, Inc., Smyrna, GA 30080.
All rights reserved. US-P-NE-RLS-2100006

Please see Important Safety Information within and refer to Patient Information for NEUPRO at the end of this brochure or at www.NEUPRO.com.